

On the Inappropriateness of Noninvasive Multidetector Computed Tomography Coronary Angiography to Trigger Coronary Revascularization

A Comparison With Invasive Angiography

Giovanna Sarno, MD, PhD,* Isabel Decraemer, BSc,† Piet K. Vanhoenacker, MD, PhD,† Bernard De Bruyne, MD, PhD,* Michalis Hamilos, MD, PhD,* Thomas Cuisset, MD,* Eric Wyffels, MD,* Jozef Bartunek, MD, PhD,* Guy R. Heyndrickx, MD, PhD,* William Wijns, MD, PhD*

Aalst, Belgium

Objectives Our purpose was to evaluate the appropriateness of multidetector computed tomography angiography (MDCTA) as an anatomical standard for decision making in patients with known or suspected coronary artery disease.

Background Although correlative studies between MDCTA and coronary angiography (CA) show good agreement, MDCTA visualizes plaque burden and calcifications well before luminal dimensions are encroached.

Methods Pressure-derived fractional flow reserve (FFR) was obtained in 81 patients (116 vessels) who underwent both CA and MDCTA. Segments were visually graded for stenosis severity as: G0 = normal, G1 = nonobstructive (<50% diameter reduction), and G2 = obstructive (≥50% diameter reduction).

Results Concordance between segmental severity scores by MDCTA and CA was good ($k = 0.74$; 95% confidence interval: 0.56 to 0.92). Diagnostic performance of MDCTA for detection of functionally significant stenosis based on FFR was low (sensitivity 79%; specificity 64%; positive likelihood ratio 2.2; negative likelihood ratio 0.3). Revascularization was considered appropriate in the presence of reduced FFR (≤ 0.75). Decision making based on MDCTA guidance would result in revascularization in the absence of ischemia in 22% of patients (18 of 81) and inappropriate deferral in 7% (6 of 81), while revascularization in the absence of ischemia would be 16% (13 of 81) and inappropriate deferral 12% (10 of 81) with decisions guided by CA. Combined evaluation of stenosis severity using both anatomy (with either CA or MDCTA) and function (with FFR) yields the highest proportion of appropriate decisions: 90% and 91%, respectively ($p = 0.0001$ vs. CA only, $p = 0.0001$ vs. MDCTA only).

Conclusions Similar to CA, anatomical assessment of coronary stenosis severity by MDCTA does not reliably predict its functional significance. (J Am Coll Cardiol Intv 2009;2:550–7) © 2009 by the American College of Cardiology Foundation

From the *Cardiovascular Center and the †Department of Radiology, OLV Hospital, Aalst, Belgium.

Manuscript received January 13, 2009; revised manuscript received March 2, 2009, accepted March 8, 2009.

Visual estimation of coronary stenosis severity during invasive coronary angiography (CA) does not predict its hemodynamic significance, even when performed and analyzed by experienced cardiologists (1,2). Yet coronary revascularization using stented angioplasty is triggered off increasingly often by anatomical imaging (3,4). This trend is likely to increase with the availability of noninvasive multidetector computed tomography coronary angiography (MDCTA). Although correlative studies between MDCTA and CA show good agreement (5,6), MDCTA visualizes plaque burden and calcifications well before luminal dimensions are encroached (7-9). As a result, the information content of MDCTA is broader than that provided by CA, which may improve the decision whether or not to proceed with revascularization.

See page 558

The purpose of this prospective study was: 1) to compare the assessment of coronary stenosis severity by MDCTA versus invasive angiography in patients with known or suspected coronary artery disease (CAD); and 2) to evaluate the appropriateness of using MDCTA as the anatomical substrate for clinical decision making with respect to revascularization. Appropriateness of revascularization was assessed by pressure-derived fractional flow reserve (FFR), a reliable invasive index of stenosis hemodynamic significance. FFR expresses the maximum achievable blood flow to the myocardium supplied by a stenotic artery as a fraction of normal maximum flow (10,11). An abnormal value ≤ 0.75 identifies stenosis that is associated with inducible ischemia (12,13) that can be relieved by stented angioplasty or bypass grafting. Operators were encouraged to base the treatment strategy on FFR results, but final individual decisions were taken on the basis of all available data, at their discretion.

Methods

Patient selection. The study population consisted of 88 patients in whom MDCTA was implemented in the diagnostic workup of chest pain, of which 7 patients were excluded because of uninterpretable or technically poor MDCTA studies. From the remaining 81 patients, 39 had stable angina (48%) and 42 (52%) had atypical chest pain. Selected patients were identified prospectively at the time of their first visit at the outpatient clinic. Exclusion criteria were as follows: atrial fibrillation, creatinine clearance below 30 ml/min, and allergy to iodinated contrast agents. Only 15 patients (19%) were known to have CAD from prior history. In the 61 (81%) patients with suspected CAD, the pre-test probability for obstructive CAD was calculated using the Duke clinical risk score, which includes type of chest pain, age, sex, and traditional risk factors (14,15). Subjects are classified as low, intermediate, or high risk. Conventional CA was performed according to local practice 26 ± 11 days after the MDCTA. Follow-up was performed

to evaluate the clinical event rates: new revascularization after the index procedure, new angina onset, myocardial infarction, and cardiac death. The study protocol was approved by the institutional ethics committee and patients gave informed consent for participation and data collection. **CA and FFR assessment.** Distal coronary pressure measurement was performed with a 0.014-inch pressure guidewire (Radi Medical Systems, Uppsala, Sweden). The wire was introduced through a 6-F guiding catheter, calibrated, advanced into the coronary artery, and positioned in the distal vessel beyond the stenosis as previously described (11). Adenosine was administered to induce maximum hyperemia, either intravenously ($140 \mu\text{g/kg/min}$) or intracoronary (at least $15 \mu\text{g}$ in the right or $20 \mu\text{g}$ in the left coronary artery) (11,16,17). FFR was calculated as the ratio of mean hyperemic distal coronary pressure measured by the pressure wire to mean aortic pressure measured by the guiding catheter. The measurement was performed twice, and FFR was taken as the average of both measurements. Interrogated coronary vessels ($n = 116$) were segmented according to the American Heart Association classification (18). Stenosis severity was graded by independent experienced observers unaware of FFR results according to the following scoring system:

- Grade 0: entirely normal vessel segment, no plaque and no wall irregularities
- Grade 1: CAD present, nonobstructive plaque, $<50\%$ diameter reduction
- Grade 2: obstructive stenosis, $\geq 50\%$ diameter reduction

MDCTA protocol. After determination of the coronary artery calcium score (24×1.2 collimation, 330-ms gantry rotation time, 0.2 pitch, 3-mm slice thickness, 120-kV tube voltage, 145-mAs tube current), MDCTA was performed using a Sensation 64 scanner (Siemens Medical Solutions, Forchheim, Germany). Scanning parameters were 64×0.6 -mm collimation, 330-ms rotation time, 3.8-mm/rotation table feed, 120-kV tube voltage, and 850-mAs tube current. Axial images were reconstructed with a slice thickness of 0.75 mm and a reconstruction increment of 0.4 mm, using a medium sharp convolution kernel at 5% intervals over the entire R-R cycle using a single segment reconstruction algorithm. Before the scan, sublingual nitroglycerine was administered, and patients with a baseline heart rate >65 beats/min were given beta-blockers (metoprolol 25 to 100 mg orally before examination, supplemented by intravenous administration as required). A

Abbreviations and Acronyms

CA = coronary angiography

CAD = coronary artery disease

FFR = fractional flow reserve

LAD = left anterior descending coronary artery

MDCTA = multidetector computed tomography angiography

PCI = percutaneous coronary intervention

Table 1. Patient Demographics

| | Suspected CAD | | | | Known CAD (n = 20) | p Value |
|--------------------------|-------------------|----------------------|-------------------------------|-----------------------|-----------------------|---------|
| | Total (n = 81) | Low Risk (n = 14) | Intermediate Risk (n = 23) | High Risk (n = 24) | | |
| Men | 60 (74) | 9 (64) | 16 (70) | 21 (88) | 14 (70) | NS |
| Age (yrs) | 62 ± 11 | 52 ± 13* | 61 ± 9 | 65 ± 9 | 67 ± 9 | <0.001 |
| BMI (kg/m ²) | 28 ± 5 | 30 ± 7 | 28 ± 4 | 28 ± 4 | 26 ± 3 | NS |
| HR (beats/min) | 71 ± 11 | 77 ± 18 | 66 ± 8 | 71 ± 10 | 74 ± 12 | NS |
| Risk factors | | | | | | |
| Smoker | 32 (40) | 0 | 10 (43) | 10 (41) | 11 (55) | NS |
| Diabetes mellitus | 14 (17) | 3 (21) | 2 (8) | 4 (17) | 4 (20) | NS |
| Hypertension | 39 (48) | 9 (64) | 9 (39) | 11 (46) | 11 (40) | NS |
| Hyperlipidemia | 41 (51) | 5 (36) | 16 (70) | 9 (38) | 10 (50) | NS |
| Family history of CAD | 40 (49) | 8 (57) | 10 (43) | 12 (50) | 11 (55) | NS |
| Calcium score (Agatston) | 518 ± 578 | 138 ± 189* | 577 ± 546 | 643 ± 749 | 521 ± 396 | 0.02 |
| Clinical presentation | | | | | | |
| Stable angina | 39 (48) | 1 (7)* | 7 (30) | 20 (83) | 10 (50) | <0.0001 |
| Atypical chest pain | 42 (52) | 13 (93)* | 16 (70) | 4 (17)† | 10 (50) | <0.0001 |
| Duke score (%) | | 15 ± 6* | 49 ± 14‡ | 89 ± 7 | — | <0.0001 |
| Medical history | | | | | | |
| Previous PCI | 7 (9) | 0 | 0 | 0 | 7 (35) | <0.0001 |
| Previous AMI | 6 (7) | 0 | 0 | 0 | 6 (30) | <0.0001 |
| MVD by CA | 28 (35) | 1 (7)* | 7 (30)† | 8 (29) | 13 (65) | <0.01 |

Values are n (%) or mean ± SD. *p < 0.01 versus intermediate-risk, high-risk, and known coronary artery disease (CAD) groups; †p < 0.05 versus known CAD group; ‡p < 0.01 versus high-risk group.
AMI = acute myocardial infarction; BMI = body mass index; CA = coronary angiography; HR = heart rate; MVD = multivessel disease; PCI = percutaneous coronary intervention.

bolus (on average 90 ml at 5 ml/s) of contrast material (iomeprol, 816.5 mg/ml [Iomeron 400, Bracco Altana Pharma, Konstanz, Germany]) was injected intravenously followed by a 50-ml saline flush. After a scan delay, the scan started automatically when the density in the aortic root reached a density value of 100 HU. All the datasets were evaluated offline on an image analysis workstation (TeraRecon Inc., San Mateo, California). MDCTA data were evaluated on axial plane source images, on curved multiplanar reconstruction, and on different maximal intensity projections.

Coronary vessel segmentation was matched with CA, and stenosis severity was graded as described in the preceding text by independent experienced observers unaware of grading results on CA.

Statistical analysis. The diagnostic performance of MDCTA and CA for the detection of functionally significant CAD as defined by FFR ≤0.75 is presented as sensitivity, specificity, positive and negative predictive values with the corresponding confidence intervals, and positive and negative likelihood ratios. Comparisons between MDCTA and FFR or CA and FFR were performed for each interrogated vessel. Continuous variables are expressed as means and standard deviation. Differences in means among groups were analyzed by a 2-sided *t* test or by 1-way analysis of variance using a Tukey-Kramer test to compare all pairs. Categorical variables are expressed as absolute numbers and percentages. Interobserver and intraobserver variability of the score

grading by MDCTA and CA and agreement between techniques were tested by kappa statistics. False negatives were defined as vessels with abnormal FFR but nonobstructive stenosis (grade 0 to 1). False positives were defined as vessels with normal FFR but obstructive stenosis (grade 2).

The proportion of patients with consistent or inconsistent treatment decisions that are in accordance with test results is shown as absolute number and percentage for each technique. Unpaired and paired comparison between proportions uses the chi-square and McNemar tests, respectively. Appropriateness by FFR refers to revascularization of hemodynamically significant stenoses and deferral of non-significant stenoses, meaning that appropriateness is based on treatment decisions that are in accordance with functional test results. Any other decision will be called inappropriate. Operators were encouraged to base the treatment strategy on FFR results, but final individual decisions were taken on the basis of all available data, at their discretion. Kaplan-Meier curves are comparing major adverse cardiac event rates between inappropriate/appropriate treatment guided by MDTCA and FFR, respectively. Comparison between the curves uses the Gehan-Breslow-Wilcoxon test.

Results

Patient demographics are shown in Table 1. Obstructive CAD (grade 2) was diagnosed in 37 (44%) patients by CA




Figure 1. Obstructive CAD by Angiography per Category of Duke Risk Score

Distribution of study population according to the Duke risk score before multi-detector computed tomography angiography and invasive evaluation. **Solid portion of bars** represent the proportion of patients shown to have obstructive coronary artery disease (CAD) by invasive coronary angiography (CA).

and in 43 (55%) patients by MDCTA, with a concordance rate of 97% per vessel and 92% per patient. The distribution of study population according to the Duke risk score and presence of obstructive CAD by CA is shown in Figure 1. The vessel of interest ($n = 116$) was the left anterior descending in 82 (71%), the left circumflex or marginal branch in 19 (15%), and the right coronary artery in 17 (14%). The number of coronary segments assessed by visual scoring of disease severity was 334. The concordance between segmental severity scores by MDCTA and CA was good ($k = 0.74$; 95% confidence interval: 0.56 to 0.92).

Appropriateness of therapeutic decisions is shown in Figure 2. There were 51 patients with appropriate treatment guided by MDCTA, of whom there were 30 patients with obstructive stenoses treated by percutaneous coronary intervention (PCI) and 21 patients with nonobstructive stenoses treated by conservative medical therapy. There were 29 patients with inappropriate treatment guided by MDCTA, of whom there were 12 patients with obstructive stenoses treated by conservative medical therapy and 17 patients with nonobstructive stenoses treated by PCI. There were 65 patients with appropriate treatment guided by FFR, of whom 28 patients with $\text{FFR} \leq 0.75$ were treated by PCI and 37 patients with $\text{FFR} > 0.75$ were treated by conservative treatment. There were 16 patients with inappropriate treatment by FFR, of whom 6 patients with $\text{FFR} \leq 0.75$ were treated by conservative medical therapy and 10 patients with $\text{FFR} > 0.75$ were treated by PCI.

Combining anatomical evaluation (with either CA or MDCTA) with functional evaluation of stenosis severity (using FFR) yields the highest proportion of appropriate

decisions: 90% and 91%, respectively ($p = 0.0001$ vs. CA alone, $p = 0.0001$ vs. MDCTA alone).

Comparison between MDCTA and FFR. The diagnostic performance of MDCTA for the detection of functionally significant stenoses ($\text{FFR} \leq 0.75$) was poor, yielding non-diagnostic positive likelihood ratios (Table 2, Fig. 3). Mean FFR was significantly ($p < 0.01$) lower between normal (grade 0) and diseased segments. However, individual data-points show a wide overlap between intermediate (grade 1) and obstructive stenoses (grade 2) with a wide range of FFR values in grade 2 stenoses. The false negative rate was 6% (7 of 116), mostly located in the mid-distal segments of the left anterior descending coronary artery (LAD) (6 of 7, 86%). The false positive rate was 26% (30 of 116), with segments located predominantly in the LAD (19 of 30, 63%) or left circumflex coronary artery (6 of 30, 20%). Agatston coronary artery calcium score (excluding 1 outlier with a score above 10.000) was not different between true positive and false negative cases but significantly higher ($p = 0.04$) in false positive (723 [451 to 994]) than in true negative cases (357 [230 to 484]).




Figure 2. (In)Appropriateness of Treatment Decisions Guided by Anatomy, Function, or Both

Percentage of appropriate and inappropriate treatment decisions based on the result of individual or combined techniques: coronary angiography (CA), multidetector computed tomography angiography (MDCTA), fractional flow reserve (FFR), CA + FFR, MDCTA + FFR. Appropriateness by FFR is not 100% because individual decisions were taken on the basis of all available data, at the operator's discretion. Combined approaches are significantly different from both CA and MDCTA. * $p < 0.001$ versus CA; $\$p < 0.001$ versus MDCTA.

| Table 2. Diagnostic Performance of MDCTA and CA for the Detection of Functionally Significant Stenosis Based on Pressure-Derived FFR Measurements | | | | | | | | | | |
|---|-------|-------|-------|-------|-----------------|--------|-----------------|--------|------|------|
| | TP, n | TN, n | FP, n | FN, n | Sensitivity*, % | 95% CI | Specificity*, % | 95% CI | LR+* | LR–* |
| MDCTA | | | | | | | | | | |
| All vessels | 26 | 53 | 30 | 7 | 79 | 61–91 | 64 | 52–74 | 2.2 | 0.3 |
| Per patient | 24 | 33 | 18 | 6 | 80 | 61–92 | 65 | 51–78 | 2.3 | 0.3 |
| CA | | | | | | | | | | |
| All vessels | 22 | 59 | 24 | 11 | 67 | 48–82 | 71 | 60–81 | 2.3 | 0.4 |
| Per patient | 20 | 38 | 13 | 10 | 67 | 47–83 | 75 | 60–86 | 2.6 | 0.4 |

*Multidetector computed tomography angiography (MDCTA) versus coronary angiography (CA), p value = NS.
CI = confidence interval; FFR = fractional flow reserve; FN = false negative; FP = false positive; LR+ = positive likelihood ratio; LR– = negative likelihood ratio; TN = true negative; TP = true positive.

Comparison between CA and FFR. The diagnostic performance of CA for the detection of functionally significant stenosis (FFR ≤ 0.75) was poor, yielding nondiagnostic positive likelihood ratios (Table 2, Fig. 4). There was a significant difference in FFR values between intermediate (grade 1) and obstructive (grade 2) stenoses, with a marked overlap of the individual datapoints and a wide range of FFR values in grade 2 stenoses. The false negative rate was 10% (11 of 116), mostly located in the mid-distal segments of the LAD (9 of 11, 82%). The false positive rate was 21% (24 of 116), with segments located predominantly in the LAD (12, 50%) or left circumflex coronary artery (5 of 24, 21%).

Clinical outcome. During the mean follow-up time of 1.8 ± 0.3 years, 12 patients had new events: 7 presenting new angina requiring hospitalization and 5 requiring revascularization. New events occurred primarily in patients with inappropriate treatment by FFR: 1 with abnormal FFR that was not revascularized, and 4 patients with nonsignificant FFR that were revascularized. The event-free survival curves

for patients with appropriate versus inappropriate treatment by MDCTA and by FFR, respectively, are shown in Figure 5. The event-free survival rate was significantly worse in the inappropriate versus the appropriate treatment group, as defined by FFR: 68 % versus 89%, respectively ($p = 0.02$).

Implications for selection of lesions for revascularization. Given the weak correlation between FFR and both MDCTA and CA, indications for revascularization purely based on anatomy will be inappropriate by FFR guidance in nearly 30% of patients (Figs. 3 and 4). Decision making based on MDCTA would result in revascularization in the absence of ischemia in 22% (18 of 81) and inappropriate deferral in 7% of cases (6 of 81). Decision making based on CA would result in revascularization in the absence of ischemia in 16% (13 of 81) and inappropriate deferral in 12% of cases (10 of 81). Decision making based on MDCTA will not result in significantly less revascularization in the absence of ischemia or inappropriate deferral than with CA. In patients inappropriately deferred by the operator despite abnormal




Figure 3. Invasive Flow Reserve Versus Stenosis Severity by Noninvasive MDCTA

Scatter plot showing a significant difference in FFR values for grade 0 ($p < 0.01$) stenoses by MDCTA compared with grade 1 and grade 2 stenoses. A wide range of FFR values is observed for grade 2 stenosis. Abbreviations as in Figure 2.





Figure 4. Invasive Flow Reserve Versus Stenosis Severity by Invasive CA

Scatter plot showing a significant difference in FFR values between grade 1 and grade 2 stenoses by CA ($p < 0.01$) with a wide range of FFR values for grade 2 stenosis. Abbreviations as in Figure 2.



FFR, reduced flow reserve was due to diffuse disease in nearly 40% of the cases, while focal stenosis eventually amenable to stented angioplasty was present in the remainder (Fig. 6).

Discussion

In patients with stable CAD, outcome-based trials (19) and guidelines (20) recommend revascularization in the presence of disabling symptoms or extensive stress-inducible ischemia. Although large cohort studies (21) have shown

statistically significant correlations between stenosis severity assessed by quantitative CA and noninvasive or invasive functional testing, variability is such that one cannot rely on the coronary angiogram for clinical decision making in the individual patient. There are many reasons why the evaluation of stenosis severity may differ whether analyzed by CA or physiology. Selective CA does not account for the extent of downstream collateral supply or the size of the myocardial segment subtended by a given coronary artery. Diffuse luminal narrowing is common and precludes the selection of a “normal” reference segment (22).

Given the performance characteristics of the current 64-slice MDCTA, we hypothesized that this new coronary imaging modality would be more accurate than CA in identifying hemodynamically significant stenoses. Indeed, MDCTA is superior to CA by providing direct visualization of plaque load (23). Opacification of the coronary lumen by intravenous contrast delivery accounts for all sources of blood supply. Tomographic imaging allows multiplanar reconstructions that render the tridimensional structure of the coronary arteries and provides a more accurate description of eccentric stenoses.

Yet the present study shows an equally poor diagnostic performance of both imaging techniques, MDCTA and CA, for the detection of functionally significant CAD. The 26% false positive rate with MDCTA was associated with increased calcium scores. Arterial wall calcifications can be present from the early stages of atherosclerosis (24). The presence of calcifications causes blooming artifacts that increase plaque volume. With MDCTA, high coronary calcium load leads to stenosis overestimation and false positive results (25).

Clinical implications. Irrespective of mechanisms, this study demonstrates that indications for revascularization based solely on anatomy will be inappropriate in 21% (24 of 116) to 26% (30 of 116) of cases. As a matter of fact, the worst clinical outcome was seen in patients with inappropriate treatment according to FFR as a standard of reference. Because acute myocardial infarction and sudden death can proceed from plaque events at mildly obstructed sites, some physicians and patients fear to defer “treatment” of nonobstructive plaque, and preventive mechanical “plaque sealing” has been advocated. However, this hypothesis has not been properly investigated thus far (26,27). Instead, stented angioplasty of nonhemodynamically significant stenoses with an FFR >0.75 was shown not to improve the patient’s prognosis or symptoms, while consuming resources unnecessarily (12,13). These results of the DEFER (Percutaneous Coronary Intervention of Functionally Nonsignificant Stenosis) study (13) were confirmed by the larger FAME (Fractional Flow Reserve Versus Angiography for Guiding Percutaneous Coronary Interventions) trial (28) showing improved outcome with reduced costs in patients with multivessel disease randomized to FFR-guided stented an-




Figure 6. Inappropriate Deferral With Decision Making Based on Anatomy

Example of a false-negative stenosis, graded as nonobstructive (grade 1) both by multidetector computed tomography angiography (A) and coronary angiography (B) while fractional flow reserve (FFR) was significantly reduced (C). Decision making driven by anatomy will result in inappropriate deferral of revascularization.

gioplasty, as opposed to intervention guided by anatomy only. These data are consistent with the COURAGE (Clinical Outcomes Utilizing Revascularization and Aggressive Drug Evaluation) trial (19) of which a recently published substudy (29) indicates that revascularization improves outcome only when substantial ischemia is present before—and reduced by—revascularization. In the present clinically oriented study, revascularization decisions were left to the operator's discretion, which was in agreement with the results of anatomic-functional evaluation in 80% of cases (much higher than the 50% reported by Meijboom et al. [30]). Noninvasive coronary imaging by MDCTA has been recommended as a screening tool for identification of patients to be referred for revascularization (31). In accordance with recently issued guidelines (32), the present study invalidates any diagnostic strategy that would not incorporate functional testing for the presence of inducible ischemia, be it performed noninvasively before catheterization or during the invasive procedure using pressure-derived FFR (33).

Reprint requests and correspondence: Dr. William Wijns, Cardiovascular Center Aalst, OLV Ziekenhuis, Moorselbaan 164, 9300 Aalst, Belgium. E-mail: William.Wijns@olvz-aalst.be.

REFERENCES

1. Fischer JJ, Samady H, McPherson JA, et al. Comparison between visual assessment and quantitative angiography versus fractional flow reserve for native coronary narrowings of moderate severity. *Am J Cardiol* 2002;90:210-5.
2. Kimball BP, Bui S, Cohen EA, Cheung PK, Lima V. Systematic bias in the reporting of angioplasty outcomes: accuracy of visual estimates of absolute lumen diameters. *Can J Cardiol* 1994;10:815-20.
3. Topol EJ, Ellis SG, Cosgrove DM, et al. Analysis of coronary PTCA practice in the United States with an insurance claims data base. *Circulation* 1993;87:1489-97.
4. Lin GA, Dudley RA, Lucas FL, Malenka DJ, Vittinghoff E, Redberg RF. Frequency of stress testing to document ischemia prior to elective percutaneous coronary intervention. *JAMA* 2008;300:1765-73.
5. Vanhoenacker PK, Heijenbroek-Kal MH, Van Heste R, et al. Diagnostic performance of multidetector CT angiography for assessment of coronary artery disease: meta-analysis. *Radiology* 2007;244:419-28.
6. Leber AW, Knez A, von Ziegler F, et al. Quantification of obstructive and nonobstructive coronary lesions by 64-slice computed tomography: a comparative study with quantitative coronary angiography and intravascular ultrasound. *J Am Coll Cardiol* 2005;46:147-54.
7. Agatston AS, Janowitz WR, Hildner FJ, Zusmer NR, Viamonte M Jr., Detrano R. Quantification of coronary artery calcium using ultrafast computed tomography. *J Am Coll Cardiol* 1990;15:827-32.
8. Schuijff JD, Pundziute G, Jukema JW, et al. Diagnostic accuracy of 64-slice multislice computed tomography in the noninvasive evaluation of significant coronary artery disease. *Am J Cardiol* 2006;15:145-8.
9. Vanhoenacker R, Wijns W, Sarno G. Multi-detector CT of the coronary arteries. Potential use of CT angiography and coronary artery calcium scoring in screening. *JBR-BTR* 2007;90:80-3.

10. Pijls NHJ, Van Gelder B, Van der Voort P, et al. Fractional flow reserve: a useful index to evaluate the influence of an epicardial coronary stenosis on myocardial blood flow. *Circulation* 1995;92:3183-93.
11. Pijls NHJ, De Bruyne B, Peels K, et al. Measurement of fractional flow reserve to assess the functional severity of coronary-artery stenoses. *N Engl J Med* 1996;334:1703-8.
12. Bech GJW, De Bruyne B, Pijls NHJ, et al. Fractional flow reserve to determine the appropriateness of angioplasty in moderate coronary stenosis: a randomized trial. *Circulation* 2001;103:2928-34.
13. Pijls NHJ, van Schaardenburgh P, Manoharan G, et al. Percutaneous coronary intervention of functionally nonsignificant stenosis: 5-year follow-up of the DEFER study. *J Am Coll Cardiol* 2007;49:2105-11.
14. Gibbons RJ, Balady GJ, Bricker JT, et al., American College of Cardiology/American Heart Association Task Force on Practice Guidelines. Committee to Update the 1997 Exercise Testing Guidelines. ACC/AHA 2002 guideline update for exercise testing: summary article. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1997 Exercise Testing Guidelines). *J Am Coll Cardiol* 2002;40:1531-40.
15. Pryor DB, Shaw L, McCants CB, et al. Value of the history and physical in identifying patients at increased risk for coronary artery disease. *Ann Intern Med* 1993;118:81-90.
16. De Bruyne B, Pijls NH, Barbato E, et al. Intracoronary and intravenous adenosine 5'-triphosphate, adenosine, papaverine, and contrast medium to assess fractional flow reserve in humans. *Circulation* 2003;107:1877-83.
17. Kern MJ, Lerman A, Bech JW, et al., American Heart Association Committee on Diagnostic and Interventional Cardiac Catheterization, Council on Clinical Cardiology. Physiological assessment of coronary artery disease in the cardiac catheterization laboratory: a scientific statement from the American Heart Association Committee on Diagnostic and Interventional Cardiac Catheterization, Council on Clinical Cardiology. *Circulation* 2006;114:1321-41.
18. Austen WG, Edwards JE, Frye RL, et al. A reporting system on patients evaluated for coronary artery disease: report of the Ad Hoc Committee for Grading of Coronary Artery Disease, Council on Cardiovascular Surgery, American Heart Association. *Circulation* 1975;51:5-40.
19. Boden WE, O'Rourke RA, Teo KK, et al., COURAGE Trial Research Group. Optimal medical therapy with or without PCI for stable coronary disease. *N Engl J Med* 2007;356:1503-16.
20. Silber S, Albertsson P, Aviles FF, et al. Guidelines for percutaneous coronary interventions: the task force for percutaneous coronary interventions of the European Society of Cardiology. *Eur Heart J* 2005;26:804-47.
21. Bartunek J, Sys SU, Heyndrickx GR, Pijls NHJ, De Bruyne B. Quantitative angiography in predicting functional significance of stenoses in unselected patient cohort. *J Am Coll Cardiol* 1995;26:328-34.
22. De Bruyne B, Hersbach F, Pijls NH, et al. Abnormal epicardial coronary resistance in patients with diffuse atherosclerosis but "normal" coronary angiography. *Circulation* 2001;104:2401-6.
23. Mollet NR, Cademartiri F, van Mieghem CA, et al. High-resolution spiral computed tomography coronary angiography in patients referred for diagnostic conventional coronary angiography. *Circulation* 2005;112:2318-23.
24. Watson KE. Pathophysiology of coronary calcifications. *J Cardiovasc Risk* 2000;7:93-7.
25. Pugliese F, Mollet NR, Runza G, et al. Diagnostic accuracy of non-invasive 64-slice CT coronary angiography in patients with stable angina pectoris. *Eur Radiol* 2006;16:575-82.
26. Mercado N, Maier W, Boersma E, et al. Clinical and angiographic outcome of patients with mild coronary lesions treated with balloon angioplasty or coronary stenting. Implications for mechanical plaque sealing. *Eur Heart J* 2003;24:541-51.
27. Meier B. Plaque sealing by coronary angioplasty. *Heart* 2004;90:1395-8.
28. Tonino AL, De Bruyne B, Pijls NH, et al. Fractional flow reserve versus angiography for guiding percutaneous coronary interventions. *N Engl J Med* 2009;360:213-24.
29. Shaw LJ, Berman DS, Maron DJ, et al., COURAGE Investigators. Optimal medical therapy with or without percutaneous coronary intervention to reduce ischemic burden: results from the Clinical Outcomes Utilizing Revascularization and Aggressive Drug Evaluation (COURAGE) trial nuclear substudy. *Circulation* 2008;117:1283-91.
30. Meijboom WB, Van Mieghem CA, van Pelt N, et al. Comprehensive assessment of coronary artery stenoses: computed tomography coronary angiography versus conventional coronary angiography and correlation with fractional flow reserve in patients with stable angina. *J Am Coll Cardiol* 2008;52:636-43.
31. Shaw LJ, Berman DS, Blumenthal RS, et al. Clinical imaging for prevention: directed strategies for improved detection of presymptomatic patients with undetected atherosclerosis-part I: clinical imaging for prevention. *J Nucl Cardiol* 2008;15:e6-19.
32. Patel MR, Dehmer GJ, Hirshfeld JW, Smith PK, Spertus JA. ACCF/SCAI/STS/AATS/AHA/ASNC 2009 appropriateness criteria for coronary revascularization. *J Am Coll Cardiol* 2009;53:530-53.
33. Wijns W, De Bruyne B, Vanhoenacker PK. What does the clinical cardiologist need from noninvasive cardiac imaging: is it time to adjust practices to meet evolving demands? *J Nucl Cardiol* 2007;14:366-70.

Key Words: computed tomography ■ coronary disease ■ revascularization.